

Supernova search at intermediate z

I. Spectroscopic analysis

G. Altavilla (1), P. Ruiz-Lapuente (1), A. Balastegui (1),
J. Méndez (1,2), S. Benetti (3), M. Irwin (4), K. Schahmanech (5),
C. Balland (6,7), R. Pain (5), N. Walton (4)

1) *Department of Astronomy, CER for Astrophysics, Particle Physics and Cosmology, University of Barcelona, Diagonal 647, E-08028, Barcelona, Spain*

2) *Isaac Newton Group of Telescopes, 38700 Santa Cruz de La Palma, Islas Canarias, Spain*

3) *INAF-Astronomical Observatory of Padova, vicolo dell'Osservatorio 5, 35122 Padova, Italy*

4) *Institute of Astronomy, University of Cambridge, Madingley Road, Cambridge. CB3 0HA, United Kingdom*

5) *LPNHE-IN2P3-CNRS-Universités Paris 6 et Paris 7, 4 place Jussieu, 75252 Paris Cedex 05 France*

6) *Institut d'Astrophysique Spatiale, Bâtiment 121, Université Paris 11, 91405 Orsay Cedex, France*

7) *Université Paris Sud, IAS-CNRS, Bâtiment 121, Orsay Cedex, France*

Abstract. We study 8 supernovae discovered as part of the International Time Programme (ITP) project “ Ω and Λ from Supernovae and the Physics of Supernova Explosions” at the European Northern Observatory (ENO). The goal of the project is to increase the sample of intermediate redshift ($0.1 < z < 0.4$) SNe Ia for testing properties of SNe Ia along z and for enlarging the sample in the Hubble diagram up to large z .

1. Intermediate z Supernovae

Though SNe Ia are extensively used in cosmological studies as distance indicators, the empirical luminosity calibration, which is crucial for any investigation, is based on the local SN sample. Though extensive comparison between high- z and the local SNeIa sample is done, the intermediate- z domain (redshifts between 0.1–0.4) is still relatively unexplored. The aim of our project is to fill up the gap between the local and the high- z sample and to provide detailed spectroscopic and photometric data to improve the knowledge of SNeIa at all z . Our observations led to the confirmation of 6 SNe Ia in the range $0.033 < z < 0.329$, one of them being a peculiar subluminal object, and 2 SNe II, both at $z \sim 0.14$ (see Table 1). The rest of the discovered objects were QSOs and active galaxies. Searches were performed with the use of the Wide Field Camera (WFC) at the 2.5m Isaac Newton Telescope (INT). Detection of the SN candidates was followed by spectroscopic identification performed with the Intermediate disper-

sion Spectrograph and Imaging System (ISIS) mounted at the 4.2m William Herschel Telescope (WHT). The confirmed SNe were then followed photometrically through various ENO telescopes.

Table 1. Summary of observations.

SN name	Observ. date	R.A.(2000.0) <i>hh mm ss</i>	δ (2000.0) <i>° ' "</i>	<i>z</i>	SN Type	Comments & best matching nearby SN
2002li	Jun. 10	15:59:03.08	+54:18:16.0	0.329	Ia	~1 week before maximum, 1990N
2002lj	Jun. 11	16:19:19.65	+53:09:54.2	0.180	Ia	~1 week past maximum, 1994D
2002lp	Jun. 10	16:40:11.45	+42:28:30.2	0.144	Ia	around maximum, 1981B
2002lq	Jun. 10	16:40:28.83	+41:14:09.1	0.269	Ia	~1 week before maximum, 1990N
2002lr	Jun. 10	22:33:12.59	+01:05:56.7	0.255	Ia	~10 days past maximum, 1994D
2002lk	Jun. 10	16:06:55.92	+55:28:18.2	0.033	Ia pec.	a few days before maximum, 1991bg
2002ln	Jun. 10	16:39:24.93	+41:47:29.0	0.138	SN II	~2 weeks past maximum, 1999em
2002lo	Jun. 11	16:39:56.42	+42:19:20.5	0.136	SN II	~1 month past maximum, 1999em

The comparison of our intermediate z SN spectra with high-signal-to-noise ratio spectra of nearby SNe has not revealed significant differences. In particular we found a good match with nearby SN spectra for all our objects (see Table 1 and Fig. 1). The only supernova still lacking a good enough match is the subluminal peculiar SN 2002lk which was caught at a premaximum phase not commonly observed in nearby subluminal SNe Ia.

Our overall results suggest that SNe Ia observed at different redshifts resemble the nearby objects in most measured aspects (spectra, light curves, kinematics). A further discussion will be given elsewhere.




Figure 1. Comparison of SN 2002lj smoothed spectrum with that of SN 1994D 7 days past maximum (Patat et al. 1996) (left panel) and with a template at the same epoch (Nobili et al. 2003) (right panel).

References

- Nobili, S., Goobar, A., Knop, R. et al. 2003, *A&A*, 404, 901
 Patat, F., Benetti, S., Cappellaro, E. et al. 1996, *MNRAS*, 278, 111